

Shiina Y,* Slavik S,* Uemura H,** McCarthy KP,** Yen Ho S.** The inferior caval vein draining into the left atrial cavity – a rare case. *Images Paediatr Cardiol* 2011;13(4):1-5

*Royal Brompton Hospital, London, UK.

**Cardiac Morphology Unit, Royal Brompton Hospital and Imperial College London, UK.

Introduction


The inferior vena cava (IVC) draining into the left atrium (LA) is exceedingly rare in the setting of the usual atrial arrangement (situs solitus).¹ We describe a patient with this unique anomaly, and its repair.

Case report

A female patient aged 9 years initially developed significant exertional dyspnoea and desaturation. These symptoms, together with clubbing of fingers and toes, prompted referral for cardiovascular assessment. On physical examination her height and weight were at the 3rd centile for age (124cm and 23kg, respectively). Oxygen saturation was 91% at rest and fell to 87% with exercise. Normal volume and symmetrical peripheral pulses were present. Abdominal examination excluded hepatosplenomegaly. Chest auscultation revealed normal first and second heart sounds with no murmurs. Symmetrical air entry was heard into both lungs without added sounds. Chest x-ray and 12-lead electrocardiogram did not show any abnormal findings.


Transthoracic echocardiogram showed situs solitus with atrio-ventricular and ventriculo-arterial concordance. The IVC appeared to drain into the LA through a restrictive tunnel, although details of this route remained unclear (Figure 1).

Figure 1: Transthoracic echocardiography before surgical repair. Right to left shunt through PFO was detected from the 4chamber view. (PFO=patent oval foramen, LV=left ventricle, RV= right ventricle).




On angiography, contrast medium passed directly from the IVC into the LA through a 2mm wide opening (Figure 2). The mean IVC blood pressure was 12mmHg and mean LA pressure was 6mmHg.

Figure 2: Angiography. A catheter was passed along the IVC directly into the LA. The IVC accepted the hepatic veins and then drained into the LA through the PFO (PFO=patent oval foramen, LA=left atrium, LV=left ventricle, IVC=inferior vena).




A dilated azygos vein (9mm) drained venous blood from the lower body into a dilated (12.4mm) superior vena cava (SVC) (Figure 3).

Figure 3: Angiography. Dilated azygos vein drained into the dilated SVC.



At surgery via a medial sternotomy, the epicardial aspect of the heart showed moderate hypoplasia of the right atrium (RA) and right ventricle. With the SVC and IVC directly cannulated, cardiopulmonary bypass and cardiac arrest were established in a standard fashion. When the right

Figure 5: Histology. ‘Septum’ was resected and stained with Masson's trichrome (x16 magnification). Toward the thin end, the myocardium was capped with thick deposition of fibrous tissue.



This patient had a smooth and uneventful post-operative course. Peripheral oxygen saturation normalized and her shortness of breath improved at 3 months follow-up. During follow up of 5 years, no evidence of arrhythmia or IVC obstruction was noted.

Discussion

Most case reports regarding such anomalous connection of the IVC to the LA indicated an abnormally large right valve of the sinus venosus, the Eustachian valve, coexisting with an atrial septal defect or PFO.^{2,3} The persistence of a large Eustachian valve in various forms is a recognized entity and, when coexisting with an interatrial defect, can lead to a right to left shunt resulting in a clinically detectable cyanosis. This right valve of the sinus venosus contributes to the inferior border of the atrial septum. When extensive, it can almost divide the RA into two chambers resulting in right-sided cor triatriatum.⁴

During early embryologic development, the right and left venous valves separate the systemic sinus venosus from the primary part of the RA. The left venous valve subsequently fuses with the developing atrial septum. As the systemic sinus venosus becomes incorporated into the RA, the right venous valve diminishes. Its rudiments usually become the crista terminalis, the valve of the IVC (Eustachian valve) and the valve of the coronary sinus (Thebesian valve).⁵ However, abnormal regression of the right valve of the sinus venosus may be present occasionally. In our case, muscularisation instead of the normal absorption possibly occurred at the portion of the right valve of the sinus venosus resulting in the large Eustachian valve diverting the blood stream from the IVC into the LA through the PFO. Considering the histological composition of the abnormal septum, it is possible that initially there was a communication between the IVC cavity and the RA as cor triatriatum dextrum but the opening became obliterated by deposition of fibrous tissue at its margins.

Regarding surgical repair, some case reports indicated that the prominent Eustachian valve was intraoperatively misinterpreted as an atrial septal defect and surgically closed, even though there was an experienced cardiac surgical team at work.⁶ Detailed imaging of the IVC drainage anatomy using angiography, CT and/or magnetic resonance imaging is therefore mandatory in cases where clinical and echocardiographic suspicion of drainage anomaly is present.

Conclusions

We present a rare congenital malformation involving anomalous drainage of the IVC into the LA. A high level of clinical suspicion in otherwise unexplained cyanosis and detailed imaging of the abnormal drainage anatomy can lead to accurate diagnosis and successful surgical treatment.

References

1. Bartram U, Fischer G, Kramer HH. Congenitally interrupted inferior vena cava without other features of the heterotaxy syndrome: report of five cases and characterization of a rare entity. *Pediatr Dev Pathol* 2008;11(4):266-273.
2. Oliver JM, Gallego P, Gonzalez A. Sinus venosus syndrome: atrial septal defect or anomalous venous connection? A multiplane transoesophageal approach. *Heart* 2002;88(6):634-638.
3. Genoni M, Jenni R, Vogt PR. Drainage of the inferior vena cava to the left atrium. *Ann Thorac Surg* 1999 ;67(2):543-545.
4. Alboliras ET, Edwards WD, Driscoll DJ. Cor triatriatum dexter: two-dimensional echocardiographic diagnosis. *J Am Coll Cardiol* 1987 Feb;9(2):334-337.
5. Gallaher ME, Sperling DR, Gwinn JL. Functional drainage of the inferior vena cava into the left atrium-three cases. *Am J Cardiol* 1963 ;12:561-566.
6. Becker A, Buss M, Sebening W. Acute inferior cardiac inflow obstruction resulting from inadvertent surgical closure of a prominent Eustachian valve mistaken for an atrial septal defect. *Pediatr Cardiol* 1999;20(2):155-157.

Contact Information

© Images in
Paediatric Cardiology
(1999-2011)

Hideki Uemura
H.Uemura@rbht.nhs.uk

